FOODTEST .hu

FOOD INTOLERANCE CENTER

Complex solution in diagnostic and treatment of food intolerance

Rapid, point-of-care test for the detection of human anti-gliadin and anti-tissue transglutaminase (tTG) antibodies in whole blood

INTENDED USE

Coeliac Monitor[™] is an immunochromatographic test designed to detect human IgA antibodies against tissue transglutaminase (tTG), the main autoantigen associated with Coeliac disease, and anti-IgA gliadin antibodies associated with the consumption of gluten. Coeliac Monitor[™] can be used to monitor the effectiveness of a gluten-free diet, which is recommended to improve symptoms of Coeliac disease.

SAMPLES

Coeliac Monitor[™] has been designed for use with fresh, non-haemolysed blood samples. The kit contains micropipettes to collect a sample from finger-prick whole blood. Plasma and serum samples can also be used.

TEST PROCEDURE

- 1. Prick a finger using the safety lancet provided.
- 2. Gently massage the finger in the direction of the puncture to obtain a drop of blood. Discard the first drop of blood and collect the next drop.
- 3. Touch the drop of blood with the micropipette.




- 4. Capillary action will fill the micropipette with the required volume of blood to the black line (15ul).
- 5. Squeeze the bulb of the micropipette to dispense the blood sample into the round window.



- 6. Wait for 60 seconds, until the blood is absorbed.
- 7. Add 2 drops of dilution buffer to round window.

8. Read the results at the end of 10 minutes.

RESULT INTERPRETATION



The Coeliac Monitor^M kit must be stored at 2 - 8°C. The expiry date is printed on the box.

SENSITIVITY AND SPECIFICITY

Evaluation	Sensitivity		Specificity	
	tīG	Gliadin	tīG	Gliadin
1	90%	58%	92%	80%
2	75%	100%	100%	99%
Overall	83%	79 %	96%	90%

ORDERING INFORMATION

Code	Description	Pack Size	Shelf-Life
CNSCM	Coeliac Monitor	5 Tests	24 months

INTERFERENCE SUBSTANCES

Interference substances that are likely to be present in blood were added to a diluent and sensitivity curves performed. None of the substances were found to affect the test performance.

CE FOR PROFESSIONAL USE ONLY

distributor: Buda Labor Ltd. HU-2049 Diósd, Álmos fejedelem str. 27. www.budalabor.hu